



Print



Close Window

[Print](#)**Submitted**

on August 08, 11:42 PM

for agu-fm12

Takuo Tsuda Paid: \$60.00, Transaction #: 177230

Credit Card Type: Visa

Credit Card Number: xxxxxxxxxxxx1035

Your abstract appears below.

Please print a copy of this page for your records.

To return to the Submission Center and check your list of submissions; click "View Submissions" in the left menu.

Proof**CONTROL ID:** 1504022**TITLE:** Sodium layer in the thermosphere (110-130 km) observed at Syowa Station (69.0S, 39.6E) in Antarctica**AUTHORS (FIRST NAME, LAST NAME):** Takuo T Tsuda¹, Xinzhao Chu², Takuji Nakamura¹, Mitsumu K Ejiri¹, Takuya Kawahara³**INSTITUTIONS (ALL):** 1. NIPR, Tachikawa, Japan.

2. University of Colorado, Boulder, CO, United States.

3. Shinshu University, Nagano, Japan.

ABSTRACT BODY:

The neutral metallic atom layers (such as sodium, iron, potassium layers) are normally distributed at a height range of 80-110 km (in the upper mesosphere and lower thermosphere). Resonance scattering lidar observations of these metallic layers have been used as an important tool for investigation of the upper mesosphere and lower thermosphere. On the other hand, ground-based observations of the neutral atmosphere at higher altitude (above 110 km) are quite limited. Recently, observations of metallic layers above 110 km up to 155 km have been reported from some stations. Such high-altitude metallic layers (so-called thermospheric metallic layers) is currently well-unknown phenomenon, and has possibility to greatly improve our understanding of the lower thermosphere.

Intensive sodium temperature lidar observations were carried out at Syowa Station (69.0S, 39.6E) in Antarctica between 2000 and 2002 as a part of JARE observations. From the observational data, we investigate thermospheric sodium layer (above 110 km). In this presentation, we will report a thermospheric sodium layer event (up to 130 km) observed on 23 September 2000. In this event, the lidar detected significant signals not only from 80-110 km but also from 110-130 km. More detailed analysis has provided the temperature and the sodium-density measurements at this height range up to 130 km. The estimated sodium density reached about tens of cm^{-3} at around 120 km. The temperature profile in the thermospheric sodium layer was fairly comparable to that from the NRLMSISE-00 model. We will show these results and discuss similarity and difference between the observed thermospheric sodium layer event and previously reported thermospheric metal layer events.

INDEX TERMS: [0355] ATMOSPHERIC COMPOSITION AND STRUCTURE / Thermosphere: composition and chemistry, [0340] ATMOSPHERIC COMPOSITION AND STRUCTURE / Middle atmosphere: composition and chemistry, [3369] ATMOSPHERIC PROCESSES / Thermospheric dynamics, [3332] ATMOSPHERIC PROCESSES / Mesospheric dynamics.

(No Image Selected)

(No Table Selected)

SPONSOR NAME: Takuo Tsuda

Additional Details

Previously Presented Material:

Contact Details

CONTACT (NAME ONLY): Takuo Tsuda

CONTACT (E-MAIL ONLY): tsuda.takuo@nipr.ac.jp

TITLE OF TEAM:

ScholarOne Abstracts® (patent #7,257,767 and #7,263,655). © [ScholarOne](#), Inc., 2012. All Rights Reserved.
ScholarOne Abstracts and ScholarOne are registered trademarks of ScholarOne, Inc.



Follow ScholarOne on Twitter

[Terms and Conditions of Use](#)

Product version number 4.0.0 (Build 56)
Build date Aug 07, 2012 12:22:26. Server tss1be0074